

Welcome to Horst Ibelgaufts'

COPE

Cytokines Online Pathfinder Encyclopaedia


Version 10.3 (July 2003)

10,300+ entries/pages; 19,700+ references; 85,000+ internal hyperlinks

including fully integrated <u>MiniCOPE Dictionaries</u> on Angiogenesis, Apoptosis, Cell lines, Chemokines, CytokineTopics, Hematology, Metalloproteinases, Virokines, Viroceptors, Virulence Factors

Please keep COPE a free service. Place a banner ad.

book logo to announce upcoming event:



COPE page structure | Subdictionaries | This version's new entries

A B C D E F G H I J K L M N O P Q R S T U V W X Y Z

REQUEST FOR HELP | COPE Reviews | Server statistics

Created, developed, and maintained by Prof Dr H Ibelgaufts
Contact information


Copyright (C) 1997-2003 by Prof H Ibelgaufts Ultra posse nemo obligatur

Horst Ibelgaufts' COPE: Cytokines Online Pathfinder Encyclopaedia

COPE Philippine Universities Library Help Initiative



Antisense and Nucleic Acid Drug Development + Cloning + Cytokine + Cytokine and Growth Factor Reviews + DNA and Cell Biology + EMBO Journal + European Cytokine Network + Genetic Engineering News + Glycobiology + Human Molecular Genetics + Hybridoma + Immunology Today + International Immunology + Journal of Computational Biology + Journal of Immunological Methods + Journal of Interferon & Cytokine Research + Journal of the National Cancer Institute + Molecular Medicine Today + Nucleic Acids Research + Nucleic Acids Symposium Series + Protein Engineering + Trends in Biochemical Sciences + Trends in Cell Biology + Trends in Microbiology + Viral Immunology + Your Journal +



COPE Homepage	Previous entry:	Next entry:	Random entry:
	CAP	<u>CAP-18</u>	<u>Placentin</u>

CAP-1

[<u>CD40-associated protein-1</u>] CAP-1 binds specifically to the cytosolic domain of <u>CD40</u> but not to <u>FAS antigen</u> (see: <u>APO-1</u>) or the two known receptors of <u>TNF</u>.

The CAP-1 protein contains a C-terminal domain with strong amino acid sequence homology with a unique domain of two putative signal transducing proteins binding to the cytosolic tail of the type 2 receptor for TNF, TRAF-1 and TRAF-2 (see: TRAF). The N-terminal portion of CAP-1 contains a RING finger motif and three zinc finger-like domains similar to those found in several regulatory proteins that interact with DNA or RNA.

The protein has been renamed <u>TRAF-3 (Tumor necrosis factor receptor-associated factor-3</u>) and is known also CRAF-1, CD40-BP, or LAP1.

date of last revision: March 2002

References: Sato T et al A novel member of the TRAF family of putative signal transducing proteins binds to the cytosolic domain of CD40. FEBS Letters 358: 113-8 (1995)

COPE page structure | Subdictionaries | This version's new entries

A B C D E F G H I J K L M N O P Q R S T U V W X Y Z

REQUEST FOR HELP | COPE Reviews | Server statistics

Created, developed, and maintained by Prof Dr H Ibelgaufts